



HAL
open science

Future summer mega-heatwave and record-breaking temperatures in a warmer France climate

Margot Bador, Laurent Terray, Julien Boé, Samuel Somot, A. Alias,
Anne-Laure Gibelin, Brigitte Dubuisson

► **To cite this version:**

Margot Bador, Laurent Terray, Julien Boé, Samuel Somot, A. Alias, et al.. Future summer mega-heatwave and record-breaking temperatures in a warmer France climate. *Environmental Research Letters*, 2017, 12 (7), pp.074025. 10.1088/1748-9326/aa751c . hal-02353113

HAL Id: hal-02353113

<https://hal.science/hal-02353113v1>

Submitted on 1 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

LETTER • OPEN ACCESS

Future summer mega-heatwave and record-breaking temperatures in a warmer France climate

To cite this article: Margot Bador *et al* 2017 *Environ. Res. Lett.* **12** 074025

View the [article online](#) for updates and enhancements.

You may also like

- [Visualization Analysis of Interrelationship between Temperature Profile and Water Distribution inside the PEFC](#)
Kaito Shigemasa, Hayate Sato, Yota Otsuki et al.
- [High-resolution simulations of the thermophysiological effects of human exposure to 100 MHz RF energy](#)
David A Nelson, Allen R Curran, Hans A Nyberg et al.
- [Can air temperature be used to project influences of climate change on stream temperature?](#)
Ivan Arismendi, Mohammad Safeeq, Jason B Dunham et al.

Environmental Research Letters



LETTER

Future summer mega-heatwave and record-breaking temperatures in a warmer France climate

OPEN ACCESS

RECEIVED

25 October 2016

REVISED

14 May 2017

ACCEPTED FOR PUBLICATION

26 May 2017

PUBLISHED

19 July 2017

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Margot Bador^{1,2,3,6}, Laurent Terray³, Julien Boé³, Samuel Somot⁴, Antoinette Alias⁴, Anne-Laure Gibelin⁵ and Brigitte Dubuisson⁵

¹ Climate Change Research Centre, School of BEES, University of New South Wales, Sydney NSW 2052, Australia

² Australian Research Council (ARC) Centre of Excellence for Climate System Science, University of New South Wales, Sydney NSW 2052, Australia

³ CERFACS/CNRS, CECI UMR 5318, Climat Environnement Couplages et Incertitudes, 42 avenue Coriolis 31057 Toulouse, France

⁴ Météo-France/CNRS, CNRM UMR 3589, Centre National de Recherches Météorologiques, 42 avenue Coriolis 31057 Toulouse, France

⁵ Météo-France, Direction de la Climatologie et des Services Climatiques, 42 avenue Coriolis 31057 Toulouse, France

⁶ Author to whom any correspondence should be addressed.

E-mail: m.bador@unsw.edu.au

Keywords: future mega-heatwave, future extreme and record-breaking temperatures, temperature amplification by soil-atmosphere interactions, EURO-CORDEX regional climate model, CMIP5 global climate model, SQR observations of daily maximum temperature, spatial clustering of extreme events

Supplementary material for this article is available [online](#)

Abstract

This study focuses on future very hot summers associated with severe heatwaves and record-breaking temperatures in France. Daily temperature observations and a pair of historical and scenario (greenhouse gas radiative concentration pathway 8.5) simulations with the high-resolution (~12.5 km) ALADIN regional climate model provide a robust framework to examine the spatial distribution of these extreme events and their 21st century evolution.

Five regions are identified with an extreme event spatial clustering algorithm applied to observed temperatures. They are used to diagnose the 21st century heatwave spatial patterns. In the 2070s, we find a simulated mega-heatwave as severe as the 2003 observed heatwave relative to its contemporaneous climate. A 20-member initial condition ensemble is used to assess the sensitivity of this future heatwave to the internal variability in the regional climate model and to pre-existing land surface conditions. Even in a much warmer and drier climate in France, late spring dry land conditions may lead to a significant amplification of summer extreme temperatures and heatwave intensity through limitations in evapotranspiration. By 2100, the increase in summer temperature maxima exhibits a range from 6 °C to almost 13 °C in the five regions in France, relative to historical maxima. These projections are comparable with the estimates given by a large number of global climate models.

1. Introduction

Recent decades have seen the repeated occurrence of hot summers in France characterized by intense heatwaves and daily record temperatures. In France and Western Europe, the 2003 summer is the warmest on record, with seasonal mean temperature anomalies of about +3.2 °C on average over France (compared to the 1981–2010 climatology; communication from Météo-France). These warm temperatures were mainly associated with two heatwaves, in June and August, the latter leading to

record values of daily minimum and maximum temperature in many regions in France (Beniston 2004, Schär *et al* 2004, Fink *et al* 2004). The August heatwave of 2003 is the most intense heatwave observed during the instrumental period over France and Central Europe and has therefore been called a mega-heatwave because of its exceptional characteristics (Miralles *et al* 2014, Fischer 2014). The 2015 summer was the second warmest on record in France since 1901 (+1.5 °C; relative to 1981–2010), with three heatwaves occurring during this summer and leading to daily temperature

extremes and new record-breaking temperatures in the core of the warm spells.

Over Western Europe, the occurrence of a heatwave is dependent on the establishment and duration of specific synoptic conditions such as the very intense atmospheric blocking in August 2003 (Fink *et al* 2004, Ogi *et al* 2005, Trigo *et al* 2005, Miralles *et al* 2014). The occurrence and persistence of this atmospheric circulation pattern are influenced by the large scale circulation and remote teleconnections (Black *et al* 2004, Cassou *et al* 2005, Ogi *et al* 2005, Black and Sutton 2006, Carril *et al* 2007). However, many studies (e.g. Black *et al* 2004, Fischer *et al* 2007, Garcia-Herrera *et al* 2010, Stéfanon *et al* 2012, Miralles *et al* 2014) have also shown that the 2003 summer extreme temperatures could not have been reached without pre-existing large negative soil moisture anomalies. Other interactions between the regional climate components can also influence heatwave temperatures, such as dust aerosols during the 2006 observed event (Nabat *et al* 2015).

Climate models suggest that human influence is expected to significantly increase the frequency, duration and intensity of heatwaves in Europe (Meehl and Tebaldi 2004, Stott *et al* 2004, Fischer and Schär 2010, Christensen *et al* 2013, Christidis *et al* 2015, Russo *et al* 2014, Lau and Nath 2014, Lemonsu *et al* 2014, Schoetter *et al* 2015, Ouzeau *et al* 2016). Over Europe, in addition to surface warming, these models also suggest a decrease in soil moisture, which can partly be explained by an increase in evapotranspiration in spring and a decrease in precipitation in summer (Collins *et al* 2013). Note that these projections are commonly assessed based on a recent reference period, often selected as the last decades of the 20th century. Very few studies have looked at the occurrence and mechanisms of future mega-heatwaves in the context of their contemporaneous climate.

Our main objective is to investigate the possible occurrence of a mega-heatwave in a much warmer mean climate as well as the relevant physical mechanisms. We focus on heatwaves as extreme as the 2003 event with respect to their contemporaneous mean climate. First, a new observed daily temperature dataset for France is used to provide homogeneous regions (or clusters) of spatially co-varying summer extreme temperature events based on a statistical clustering technique. Then, an historical and 21st century climate simulation performed with the ALADIN regional climate model (at ~ 12 km spatial resolution) is used to identify the occurrence of future record-breaking temperatures and related mega-heatwaves. We find a simulated mega-heatwave in the second part of the century (in summer 2075) that is as extreme as that of 2003 in terms of anomalous extreme temperatures. A 20-member initial condition ensemble initialized one month before the heatwave onset is used to examine a possible influence of land-atmosphere interactions on surface extreme tempera-

ture amplification. We focus on the influence of pre-existing soil moisture conditions and related feedbacks which are known to play a significant role on the heatwave temperatures under current conditions. Finally, we use observed current record-breaking temperatures and their projected changes to provide a rough estimate of the summer daily temperature maxima that could be reached by 2100.

2. Data and method

2.1. Observed and simulated daily maximum temperatures

Observed daily maximum temperatures (T_{\max}) over France are taken from the SQR (Séries Quotidiennes de Référence, Reference Daily Series in English) dataset developed by Météo-France. The monthly series were carefully selected and homogenized from 1950 to 2012 (Gibelin *et al* 2014). From this first selection, non-homogenized T_{\max} were extracted, with a threshold of maximum number and magnitude of breaks detected when homogenizing the monthly series (see Dubuisson and Moisselin 2006 for more details and figure S1 stacks.iop.org/ERL/12/074025/mmedia for the spatio-temporal distribution of the stations).

We consider two simulations of the ALADIN (version 5.3) regional climate model (RCM; Colin *et al* 2010, Trambly *et al* 2013) at a very high-resolution (0.11° or ~ 12.5 km): one historical (ALADIN-HIST, with observed natural and anthropogenic forcings, 1950–2005) and the continuing future simulation (ALADIN-SCEN, 2006–2100) under the business-as-usual scenarios of green house gases (GHG) and aerosols emissions, the Representative Concentration Pathways 8.5 (RCP8.5; van Vuuren *et al* 2011). ALADIN-HIST and ALADIN-SCEN share the same vegetation and land cover. These two simulations are part of the World Climate Research Program Coordinated Regional Downscaling Experiment for Europe (EURO-CORDEX, Jacob *et al* 2014). The lateral boundary conditions and the initial conditions prescribed to ALADIN are provided by the CNRM-CM5 global climate model (GCM; Voldoire *et al* 2012). Note that the driving CNRM-CM5 GCM and the driven ALADIN RCM have very similar physical parameterizations ensuring physical consistency.

We also use a large (20 members) initial condition ensemble (ALADIN-LE) of simulations of the 2075 summer. The ensemble simulations use the EURO-CORDEX domain and cover the full 92-day period of the summer months (June, July and August; JJA). All ensemble members have exactly the same lateral and initial conditions as those of the ALADIN-SCEN simulation (the first member of the ensemble). A small random offset ($\sim 10^{-4}$ ppm) to the initial CO_2 concentration was added to each ensemble member. This very small perturbation serves to create spread among the ensemble members as internally-generated

variability increases with time (see Sanchez-Gomez and Somot 2016 for more details on ALADIN internal variability over a Euro-Mediterranean domain).

Finally, an ensemble of historical and RCP8.5 simulations from 29 global climate models (one simulation per model; see supplementary table for the list of models) of the Coupled Model Intercomparison Project Phase 5 (CMIP5; Taylor *et al* 2012) is also used to estimate an uncertainty range for the summer record maximum value in 2100, as defined below.

2.2. Summer record maximum value

On a given day, a record is broken at the n th year after the initialization when T_{\max} (year = n) is greater than the values of all previous years on that day. Records are independently estimated for each JJA calendar day and for each station (or model grid point). In this paper we are interested in the number of records broken in summer as well as the value of the record, as developed below.

We first focus on the yearly evolution of the total number of records broken in summer over France (i.e. each year the number of broken records is simply the total number of records summed over all JJA days and stations). This record evolution is shown under its normalized form, meaning that we compare this number with respect to that of a stationary climate (1 for all years) which is assumed to be consistent with the beginning of the observations (Bador *et al* 2016; see also Supporting Information S1 for more details).

We also focus on the actual value of a record-breaking temperature, and more specifically on the summer record maximum value (referred to as the record maximum value thereafter). The record maximum value in year X corresponds to the maximum value of T_{\max} over all the summer days from the initial year to year X. In a given year, record-breaking temperatures are a necessary but not sufficient condition for a record maximum value to be broken. A record maximum value is established if across each JJA day at least one of the record-breaking temperatures exceeds the previous record maximum value.

2.3. Spatial clustering of extreme events


To identify homogeneous regions in France in terms of summer extreme temperatures we use a clustering algorithm originating from multivariate extreme value theory (Coles *et al* 2001, Beirlant *et al* 2004, de Haan and Ferreira 2006, Resnick 2007). This technique is suitable to extract spatial dependencies from series of seasonal maxima. Stations with a similar interannual evolution of summer T_{\max} maximum are gathered together in K clusters. Note that no geographical information is used to cluster the stations. Instead, we use a statistical distance measuring the proximity of two stations with respect to their maxima (Bernard *et al* 2013 and Bador *et al* 2015; see supporting information S2 for more details).

The clustering algorithm is applied to the SQR observations over the period with maximum and consistent station data availability (1980–2010, see figure S1). The algorithm identifies five well defined geographical regions: South-West France (SW), Eastern France (EA), Brittany (BR), Northern France (NO) and the Mediterranean region (ME). These regions can be used to characterize the typical spatial extent of record-breaking temperatures and heatwaves (figure 1). ME and SW are clearly separated due to competing influences from the Mediterranean Sea and the Atlantic. The sensitivity of the results to the prescribed number of clusters (K) is documented in Supporting Information (figure S2). It is interesting to note that the ME region shows the strongest cluster strength for different K number of clusters. This highlights the very particular summer climate of this small region (in terms of temperature extremes) largely influenced by the Mediterranean Sea. It also demonstrates the advantage of using a clustering of extreme events as a geographical selection of 5 regions in France would probably result in a larger ME cluster, with more stations taken around the Mediterranean edge and further inland. All five regions have observed current record maximum values greater or equal to 42 °C, with the warmest being ME with 42.7 °C. We use the ALADIN model closest grid point to every station (see figure S3) to look at the changes in record-breaking temperatures occurring during the 21st century in the five regions in France.

3. Results

Figure 2 (top panel) shows the 21st century evolution of the total number of records broken in summer over France as simulated by ALADIN-SCEN. Compared to the first half of the 21st century the second half features a higher frequency of large amplitude record-breaking events and a robust emergence of the anthropogenic signal (Elguindi *et al* 2012, Bador *et al* 2016). It comprises three exceptional heatwaves in 2075, 2097 and 2099. The simulated heatwave in 2075 is characterized by a normalized number of records around 20, which means that 20 times more records are broken this summer than expected that far in a stationary climate time series assumed to be consistent with the summers of the first decades of observations. It is also associated with many new record maximum values, in particular in the EA and NO regions (figure 2, bottom panel). Despite the warming trend due to GHG concentration increase, many of the newly established record maximum values remain as such until 2100. This is particularly the case in NO and EA. For the SW, BR and ME regions, the record maximum values in 2100 are reached after year 2095 (see figure S3).

We now focus on the 2075 summer heatwave and ask how exceptional this event is in the context of its contemporaneous climate. To get some insight on the




extreme character of the simulated 2075 event, we compare it with the observed 2003 summer heatwave. Figure 3 shows the changes in record maximum values between 2075 and 2074, and 2003 and 2002, for the simulated and observed heatwaves, respectively. Note that only warm years can change the record maximum values as by definition this metric includes the memory of all previous summers up to the considered year. Both observed 2002 and simulated 2074 summers are within the range of their contemporaneous climate and cause almost no change in record maximum value in France (figure S1 and figure 2, respectively). Hence, looking at the difference to the previous year 2002 (2074) does not artificially influence the magnitude of the 2003 (2075) heatwave in terms of broken records. The spatial extent of the 2075 heatwave is slightly smaller than that of the 2003 heatwave, whereas its amplitude is larger, in particular in North-eastern France (EA and, to a lesser extent, NO). The 2075 heatwave features new record maximum values that exceed previous records by 1°C–6°C whereas the range for the 2003 heatwave is between 1° and 4°C.

We now examine in more detail the precise timing the 2075 heatwave. The heatwave duration is defined using the heatwave index as in Schoetter *et al* (2015), which is estimated in the driving CNRM-CM5 simulation over Europe (35°N–60°N; 10°W–30°E). Based on this index a heatwave is detected if at least 30% of the domain is affected for at least 3 consecutive

days by T_{\max} exceeding the 98th percentile of the T_{\max} distribution with respect to a base period (May to October months from 1979 to 2008). Figures 4(a) and (b) shows the evolution of T_{\max} anomalies (relative to late 21st century climate) averaged over Europe and France during summer 2075 in the CNRM-CM5 simulation (thick red line). The heatwave onset begins after 35 days (day 187 of the year) and becomes a real heatwave after 40 days (day 192; red shading, figure 4(a)). In France this heatwave is composed of two peaks (figure 4(b)).


Then we use the ALADIN-LE simulations (thin colored lines, figures 4(a) and (b)) to describe the 2075 heatwave and investigate the associated mechanisms. Over Europe, the ensemble spread (i.e. dispersion between the members) is small for the first 20 days and mean T_{\max} anomalies are small (less than 2°C; figure 4(a)). The ALADIN-LE spread in mean T_{\max} anomalies varies with time and is characterized by 10-to-20 day periods with large internal variability, in particular over France during the first peak of the heatwave (figure 4(b)). The first 15–20 days of the heatwave are characterized by a large spread that increases and then slowly saturates with time over Europe and France. One member completely fails to reproduce the heatwave (it actually simulates a cooling during the first 15 days of the event) before merging back with the rest of the ensemble 10 days before the end of the heatwave (solid orange line, figure 4(a)). Despite the strong



constraint due to lateral and sea surface boundary conditions, the ALADIN model thus develops a high level of internal variability within the regional domain (Sanchez-Gomez and Somot 2016). This internal variability has a stronger influence than the large-scale and sea surface forcing from the CNRM-CM5 driving model in one case out of 20. The other 19 members all simulate a heatwave albeit with very different amplitudes for both Europe and France. The T_{\max} anomaly difference between the two extreme members (among the 19 reproducing the heatwave)

can reach 2.5°C–8.3°C at any time during the heatwave when averaged over Europe (figure 4(a)). The temperature spread can even reach higher values (13.3°C) when averaged over France (figure 4(b)).


The atmospheric circulation forcing the 2075 heatwave is typical for western European heatwaves (Cassou *et al* 2005). The positive temperature anomalies maintained by a blocking anticyclone over Europe are initially due to the advection of hot air by a strong south-westerly flow (see figure S5). The 19 members of ALADIN-LE that simulate the heatwave



have a large-scale atmospheric pattern (based on daily sea level pressure—SLP; figure 4(c)) that is similar to the CNRM-CM5 driving model as soon as the heatwave begins. Indeed, the spatial pattern correlations between the 19 members of ALADIN-LE and CNRM-CM5 daily SLP range from 0.84 to 0.94 in average during the first 5 days of the heatwave (days 192–196; figure 4(c)). The similarity of the large-scale atmospheric circulation pattern among the 19 members suggests that the atmospheric circulation is likely not the main driver of the different heatwave intensity within the ALADIN-LE. The remaining member (which simulates a cooling; solid orange line) has a weaker correlation of 0.66 in average during the first 5 days of the heatwave (days 192–196; figure 4(c)). This suggests that this particular member does not

simulate a heatwave because it does not follow the large-scale pattern of the driving CNRM-CM5 model. Interestingly, the SLP pattern correlations between ALADIN-LE and CNRM-CM5 exhibit a much larger spread just before the onset of the heatwave (days 180–192). All 20 ALADIN-LE member correlations are scattered throughout the 0.43–0.92 range. This is a nice illustration of the intermittent character of the internal variability generated by regional climate models (Giorgi and Bi 2000, Alexandru *et al* 2007, Lucas-Picher *et al* 2008, Sanchez-Gomez and Somot 2016).

We take advantage of the large temperature spread of the ALADIN-LE ensemble during the heatwave to identify the mechanisms that can lead to an amplification of its temperatures. Previous



studies (e.g. Black *et al* 2004, Fischer *et al* 2007, Garcia-Herrera *et al* 2010, Stéfanon *et al* 2012, Miralles *et al* 2014) focused on present-day climate have documented the importance of pre-existing soil moisture conditions on the amplification of extreme temperature during heatwaves. Here we follow the same track to investigate if these mechanisms can also enhance the amplitude of an extreme heatwave in a mean climate that is much warmer and drier than the present-day climate. For this additional analysis, the

member without heatwave is removed from the ALADIN-LE ensemble then reduced to a 19-member ensemble.

We isolate two groups among the members using a composite analysis based on the mean soil moisture anomaly (relative to late 21st century climate) before the onset of the heatwave. Note that we use the total soil moisture integrated over a depth that varies spatially but typically corresponds to values of about 1.5m in France. The first group includes the five

members with the largest positive soil moisture anomaly averaged over the domain and over a 5-day period just before the heatwave onset (days 184–188), whereas the second one includes the five with the largest negative values. We apply this analysis to both Europe and the most impacted region of France (EA; figures 4(d) and (e)). The two groups are significantly different in terms of the mean Europe and EA T_{\max} anomalies during the first days of the heatwave, whereas they become indistinguishable later on. The temperature difference between the two groups varies between 1.9 and 6.1 °C throughout the first 10 days of the heatwave in average over both geographical domains. This suggests that pre-existing soil moisture anomalies can strongly modulate the heatwave intensity.


Soil moisture is an important climatic control on the partitioning between sensible and latent heat flux, thereby modulating air temperature. We now detail the mechanism leading to the heatwave temperature amplification and the regional spatial contrast over France. We focus on the first 10 days of the heatwave (days 192–201) when there is a significant relation between the temperature spread and the pre-existing soil moisture conditions (figures 4(d) and (e)). First, we investigate the relationship between the mean anomalies in soil moisture before the heatwave and the mean anomalies in evapotranspiration during the first days of the heatwave in the ALADIN-LE 19-member ensemble (figure 5, left column). In all regions, evapotranspiration increases with soil moisture content as indicated by significant inter-member correlations (p -values < 0.01 for all regions except NO with p -values = 0.015). To confirm a control of T_{\max} by the pre-existing soil conditions we now quantify the relationship between evapotranspiration and T_{\max} during the first days of the heatwave (figure 5, middle column). T_{\max} anomalies are mainly linked to evapotranspiration anomalies in SW and ME (inter-member correlation of -0.90 and -0.86 , respectively, with p -values < 0.01). EA and NO regions also show a strong relationship with a correlation of -0.79 and -0.64 (p -values < 0.01) whereas no significant relationship is found for BR (correlation of -0.41). Mean T_{\max} anomalies are larger in EA and NO than in ME and SW by roughly 3 °C. The origin of the heatwave pattern geographical distribution could be traced back to the spatial distribution of total soil moisture and a higher sensitivity of evapotranspiration to soil moisture changes when the soil moisture content is already low (Planton *et al* 2005). EA and NO are the two regions the most impacted by the heatwave and these two regions exhibit drier conditions than the other regions before the heatwave (days 184–188; figure 5, right column).

The results of the ALADIN-LE simulations demonstrate that very severe heatwaves with respect to their contemporaneous mean climate can occur in

a much warmer and drier French regional climate such as projected under the RCP8.5 scenario. They also suggest that mechanisms similar as those operating during present-day heatwave amplification could still be important. The future occurrence of such mega-heatwaves is expected to lead to a large increase in T_{\max} that human and ecosystems will have to cope with. Here we quantify the record maximum values that could possibly be reached in France at the end of the 21st century. We use a simple approach based on an anomaly framework where we combine observed current record maximum values with future changes as simulated by the ALADIN-SCEN simulation. By the end of the observed period, ALADIN slightly underestimates the observed record maximum values (see the right side of figure 2 and see also table 1). This might not be associated with a cold bias. Indeed the probability for a heatwave as intense as observed in 2003 is very low under present-day conditions (Schär *et al* 2004, Trigo *et al* 2005). Schoetter *et al* (2015) and Russo *et al* (2014) show that none of the CMIP5 models reproduce an event of such an extraordinary nature over the historical period. By 2100, ALADIN-SCEN indicates mean regional changes ranging from 4.4 °C–6.6 °C (averaged over NO and EA, respectively). Regional maximum increases show that individual locations can experience even greater changes: 6.6 °C, 7.7 °C, 7.7 °C, 9.6 °C and 9.9 °C for BR, NO, ME, SW and EA, respectively.

A complete assessment of the different sources of uncertainty and their contribution to the record maximum values in France is beyond the scope of our study as it would require a very large ensemble of simulations with many RCMs (and many members per model), many driving GCMs and different GHG scenarios. However, the ALADIN-LE ensemble can be used to give a qualitative estimate of the influence of internal variability (within the RCM domain) on the 2075 heatwave and associated record maximum values in 2100 (table 1). For instance, regional maximum changes 4.5 °C and 3 °C higher than simulated by ALADIN-SCEN are found in NO and EA (respectively), the two regions the most impacted by the heatwave. Finally, the combined use of ALADIN-SCEN and ALADIN-LE simulations shows that the model can simulate a change in record maximum value up to 12.9 °C in France (table 1).

It is possible to give an estimate of the model uncertainties around these projections using the CMIP5 multi-model ensemble. We use the CMIP5 ensemble (29 models) to benefit from a larger range of models compared to the EURO-CORDEX ensemble, which is only composed of a small number of driving GCMs and driven RCMs. We focus on the record maximum value in France as the CMIP5 models have a too coarse spatial resolution to attribute meaningful record values at regional and sub-regional scales. The



CMIP5 median and inter-quartile range of record maximum values in France are 11.8°C and [9°C–13.2°C], respectively. Hence, the 12.9°C increase in record maximum value projected by ALADIN appears

to be within the 3rd quartile of the CMIP5 ensemble. The ALADIN model thus simulates regional changes comparable to the estimates of the global climate models.

Table 1. Regional observed (SQR station dataset) and simulated (ALADIN-HIST simulation) current summer record maximum values ($^{\circ}\text{C}$) and their simulated changes and uncertainties given by the ALADIN and CMIP5 simulations.

summer record maximum value ($^{\circ}\text{C}$)						
in 2005		changes in 2100 compared to 2005				
SQR observations	ALADIN-HIST simulations	ALADIN-SCEN simulation		ALADIN-SCEN + ALADIN-LE simulations	CMIP5 ensemble median [inter-quartile]	
regional max	regional max	regional mean	regional max	regional max	France max	
ME	42.7	41.8	5.8	7.7	7.7	
NO	42.2	39.8	4.4	7.7	12.2	
BR	42.0	40.4	4.5	6.6	6.6	11.8 [9.0, 13.2]
EA	42.4	40.0	6.6	9.9	12.9	
SW	42.0	42.6	6.2	9.6	9.6	

4. Summary and discussion

By applying a spatial clustering methodology to temperature observations, we have identified 5 homogeneous regions in France with regard to summer temperature extremes. Based on a future simulation of the ALADIN RCM under the RCP8.5 scenario, we have shown that a summer heatwave at least as severe relative to its contemporaneous climate as the observed 2003 heatwave could occur in France in the late 21st century. This simulated heatwave has a distinct spatial pattern with the largest anomalies occurring in Northern and Eastern France.

We have then used a large ensemble (20 members) of 3-month ALADIN simulations to investigate the physical processes involved in shaping the heatwave intensity. While all ensemble members are constrained to follow the same large-scale circulation pattern driving the heatwave, the heatwave intensity and associated temperature anomalies and records differ by a large amount across the ensemble. In one case, the internal variability within the RCM domain is even able to reverse the sign of the temperature anomaly. By initializing the ensemble one month before the heatwave onset, we generate different land surface conditions across the different members simulating the heatwave. Thanks to a composite analysis based on pre-existing soil moisture content, we have then investigated the influence of soil moisture conditions and related feedbacks on the heatwave magnitude and spatial pattern. In agreement with similar studies of observed present-day heatwaves, the results show that the regions with the driest conditions before the heatwave experience large temperature anomalies and a higher number of record-breaking temperatures during the heatwave. The results also show that regional heatwave temperature anomalies can vary by several degrees due to different soil water conditions prior to the heatwave.

Finally, we have provided a qualitative estimate of the summer record maximum value that could be reached in 2100 under RCP8.5. Based on an anomaly framework, we have combined observed current

record maximum value (above 42°C) with 21st century changes simulated with ALADIN (from 6°C to almost 13°C). This ALADIN simulated change appears to belong to the inter-quartile range of the CMIP5 ensemble that indicates a median value of 11.8°C . This last estimate can be compared to changes in mean summer temperature in France that have been estimated to be $\sim 6^{\circ}\text{C}$ from an ensemble of CMIP5 models under RCP8.5 (Terray and Boé 2013). In France, the change in summer maxima of daily maximum temperatures is thus expected to be twice as large as the mean summer change.

Results indicate that record maximum value in France could easily exceed 50°C by the end of the 21st century. These extreme temperatures are experienced in desert regions, which question how realistic such projected extreme temperatures could be in France. Apart from the potential limitations associated with both GCM and RCM physics, the vegetation could be further investigated as it also controls the heatwave magnitude (Stéfanon *et al* 2012) and will experience changes with global warming. Besides, the temperature amplification by land-atmosphere interactions may not be representative of all types of heatwave in mid-latitude locations and all seasons. Here we selected summer heatwaves having a strong influence on daily maximum temperatures but heatwaves can also strongly impact night-time temperatures with serious consequences for human health.

Acknowledgments

We thank the anonymous reviewers for their helpful and constructive comments, which have improved this manuscript. MB is supported by the Australian Research Council (ARC) Discovery Grant DPI160103439 and Centre of Excellence for Climate Systems Science Grant CE110001028. This work is also supported by EDF and by the French National Research Agency and its program ‘Investissement d’avenir’ under the grant ANR-11-RSNR-0021. We acknowledge the modeling groups for access to the CMIP5 models

and data. The analyses and graphics have been done using the NCAR Command Language (NCL 2013) and the R package 'ClusterMax' available on Philippe Naveau personal web page. We also thank Robert Schoetter for the heatwave index calculation.

References

- Alexandru A, de Elia R and Laprise R 2007 Internal variability in regional climate downscaling at the seasonal scale *Mon. Weather Rev.* **135** 3221–38
- Bador M, Naveau P, Gilleland E, Castellà M and Arivelo T 2015 Spatial clustering of summer temperature maxima from the CNRM-CM5 climate model ensembles and E-OBS over Europe *Weather Clim. Extr.* **9** 17–24
- Bador M, Terray L and Boé J 2016 Emergence of human influence on summer record-breaking temperatures over Europe *Geophys. Res. Lett.* **43** 404–12
- Beirlant J, Goegebeur Y, Segers J and Teugels J 2004 *Statistics of extremes: Theory and applications* (New York: John Wiley & Sons)
- Benestad R E 2003 How often can we expect a record event? *Climate Res.* **25** 3–13
- Beniston M 2004 The 2003 heatwave in Europe: A shape of things to come? An analysis based on Swiss climatological data and model simulations *Geophys. Res. Lett.* **31** L02202
- Bernard E, Naveau P, Vrac M and Mestre O 2013 Clustering of Maxima: Spatial Dependencies among Heavy Rainfall in France *J. Clim.* **26** 7929–37
- Black E, Blackburn M, Hoskins B and Methven J 2004 Factors contributing to the summer 2003 European heatwave *Weather* **59** 217–23
- Black E and Sutton R 2006 The influence of oceanic conditions on the hot European summer of 2003 *Clim. Dyn.* **28** 53–66
- Carril A F, Gualdi S, Cherchi A and Navarra A 2007 Heatwaves in Europe: areas of homogeneous variability and links with the regional to large-scale atmospheric and SSTs anomalies *Clim. Dyn.* **30** 77–98
- Cassou C, Terray L and Phillips A S 2005 Tropical Atlantic Influence on European Heatwaves *J. Clim.* **18** 2805–11
- Christensen J H *et al* 2013 Climate phenomena and their relevance for future regional climate change *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge: Cambridge University Press) ch 14
- Christidis N, Jones G S and Stott P A 2015 Dramatically increasing chance of extremely hot summers since the 2003 European heatwave *Nat. Clim. Change* **5** 46–50
- Coles S, Bawa J, Trenner L and Dorazio P 2001 *An introduction to statistical modeling of extreme values* (Vol. 208) (London: Springer)
- Colin J, Déqué M, Radu R, and Somot S 2010 Sensitivity study of heavy precipitation in Limited Area Model climate simulations: Influence of the size of the domain and the use of the spectral nudging technique *Tellus, Series A: Dynamic Meteorology and Oceanography* **62** 591–604
- Collins M *et al* 2013 Long-term climate change: projections, commitments and irreversibility *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge, NY: Cambridge University Press)
- de Haan L and Ferreira A 2006 Extreme Value Theory, An Introduction *Springer Series in Operations Research and Financial Engineering* (New York: Springer-Verlag)
- Dubuisson B and Moisselin J 2006 Evolution des extrêmes climatiques en France à partir des séries observées *La Houille Blanche* **6** 42–7
- Elguindi N, Rauscher S a and Giorgi F 2012 Historical and future changes in maximum and minimum temperature records over Europe *Clim. Change* **117** 415–31
- Fink A H, Bruecher T, Krueger A, Leckebusch G C, Pinto J G and Ulbrich U 2004 The 2003 European summer heatwaves and drought-synoptic diagnosis and impacts *Weather* **59** 209–16
- Fischer E M, Seneviratne S I, Vidale P L, Lüthi D and Schär C 2007 Soil moisture—atmosphere interactions during the 2003 European summer heatwave *J. Clim.* **20** 5081–99
- Fischer E M and Schär C 2010 Consistent geographical patterns of changes in high-impact European heatwaves *Nat. Geosci.* **3** 398–403
- Fischer E M 2014 Climate science: Autopsy of two mega-heatwaves *Nat. Geosci.* **7** 332–3
- García-Herrera R, Diaz J, Trigo R M, Luterbacher J and Fischer E M 2010 A review of the European summer heatwave of 2003 *Crit. Rev. Environ. Sci. Technol.* **40** 267–306
- Gibelin A L, Dubuisson B, Corre L, Deaux N, Jourdain S, Laval L, Piquemal J M, Mestre O, Dennetière D, Desmidt S and Tamburini A 2014 Évolution de la température en France depuis les années 1950: Constitution d'un nouveau jeu de séries homogénéisées de référence *La Météorologie* **87** 45–53
- Giorgi F and Bi X 2000 A study of internal variability of a regional climate model *J. Geophys. Res.* **105** 29503
- Jacob D, Petersen J, Eggert B, Alias A, Christensen O B, Bouwer L M and Georgopoulou E 2014 EURO-CORDEX: new high-resolution climate change projections for European impact research *Reg. Environ. Change* **14** 563–78
- Lau N-C and Nath M J 2014 Model simulation and projection of European heatwaves in present-day and future climates *J. Clim.* **27** 3713–30
- Lemonsu A, Beaulant A, Somot S and Masson V 2014 Evolution of heat wave occurrence over the Paris basin (France) in the 21st century *Clim. Res.* **61** 75–91
- Lucas-Picher P, Caya D, de Elia R and Laprise R 2008 Investigation of regional climate models' internal variability with a ten-member ensemble of 10-year simulations over a large domain *Clim. Dyn.* 927–40
- Meehl G A and Tebaldi C 2004 More intense, more frequent, and longer lasting heatwaves in the 21st century *Science* **305** 994–7
- Meehl G A, Tebaldi C, Walton G, Easterling D and McDaniel L 2009 Relative increase of record high maximum temperatures compared to record low minimum temperatures in the US *Geophys. Res. Lett.* **36** L23701
- Miralles D G, Teuling A J and van Heerwaarden C C 2014 Mega-heatwave temperatures due to combined soil desiccation and atmospheric heat accumulation *Nat. Geosci. Lett.* **7** 345–9
- Nabat P, Somot S, Mallet M, Sevault F, Chiacchio M and Wild M 2015 Direct and semi-direct aerosol radiative effect on the Mediterranean climate variability using a coupled regional climate system model *Clim. Dyn.* **44** 1127–55
- Ogi M, Yamazaki K and Tachibana Y 2005 The summer northern annular mode and abnormal summer weather in 2003 *Geophys. Res. Lett.* **32** 1–4
- Ouzeau G, Soubeyroux J M, Schneider M, Vautard R and Planton S 2016 Heat waves analysis over France in present and future climate: application of a new method on the EURO-CORDEX ensemble *Clim. Serv.* **4** 1–12
- Planton S, Déqué M, Douville H and Spagnoli B 2005 *Impact of Climate Warming on the Hydrological Cycle* **337** 193–202
- Resnick S I 2007 *Heavy-Tail Phenomena: Probabilistic and Statistical Modeling* (New York: Springer Science & Business Media)
- Russo S, Dosio A, Graversen R G, Sillmann J, Carrao H, Dunbar M B and Vogt J V 2014 Magnitude of extreme heatwaves in present climate and their projection in a warming world *J. Geophys. Res.: Atmospheres* **119** 1–13

- Sanchez-Gomez E and Somot S 2016 Internal variability in a Mediterranean regional climate model: from climate to synoptic scales *Clim. Dyn.* (accepted)
- Schär C, Vidale P L, Lüthi D, Frei C, Häberli C, Liniger M A and Appenzeller C 2004 The role of increasing temperature variability in European summer heatwaves *Nature* **427** 332–6
- Schoetter R, Cattiaux J and Douville H 2015 Changes of western European heat wave characteristics projected by the CMIP5 ensemble *Clim. Dyn.* **45** 1601–16
- Stéfanon M, Drobinski P, D'Andrea F and De Noblet-Ducoudré N 2012 Effects of interactive vegetation phenology on the 2003 summer heatwaves *J. Geophys. Res.: Atmospheres* **117** 1–15
- Stott P A, Stone D A and Allen M R 2004 Human contribution to the European heatwave of 2003 *Nature* **432** 610–4
- Taylor K E, Stouffer R J and Meehl G A 2012 An overview of CMIP5 and the experiment design *Bull. Am. Meteorol. Soc.* **93** 485–98
- Terray L and Boé J 2013 Quantifying 21st-century France climate change and related uncertainties *C. R. Geosci.* **345** 136–49
- Tramblay Y, Ruelland D, Somot S, Bouaicha R and Servat E 2013 High-resolution med-CORDEX regional climate model simulations for hydrological impact studies: a first evaluation of the ALADIN-climate model in morocco *Hydrol. Earth Syst. Sci.* **17** 3721
- Trigo R M, García-Herrera R, Diaz J, Trigo I F and Valente M A 2005 How exceptional was the early August 2003 heatwave in France? *Geophys. Res. Lett.* **32** L10701
- Vautard R *et al* 2013 The simulation of European heat waves from an ensemble of regional climate models within the EURO-CORDEX project *Clim. Dyn.* **41** 2555–75
- Van Vuuren D P, Edmonds J, Kainuma M, Riahi K, Thomson A, Hibbard K and Masui T 2011 The representative concentration pathways: an overview *Clim. Change* **109** 5
- Voldoire A, Sanchez-Gomez E, Salas y Méria D, Decharme B, Cassou C, Sénési S and Chauvin F 2012 The CNRM-CM5.1 global climate model: description and basic evaluation *Clim. Dyn.* **40** 2091–121